



Age	Required	Recommended
Newborn	Hepatitis B #1	
1 month		
2 month	Hepatitis B #2	Rota #1
	DTaP (diphtheria, tetanus, and pertussis) #1	
	IPV (Polio) #1	
	Hib (H. influenza type B) #1	
	Pneumococcal #1	
4 month	DTaP (diphtheria, tetanus, and pertussis) #2	Rota #2
	IPV (Polio) #2	
	Hib (H. influenza type B) #2	
	Pneumococcal #2	
6 month	Hepatitis B #3	Rota #3
	DTaP (diphtheria, tetanus, and pertussis) #3	
	IPV (Polio) #3	
	Hib (H. influenza type B) #3	
	Pneumococcal #3	
9 month		
12 month	Pneumococcal #4	
	MMR (Measles, Rubella and Mumps) #1	
	Hepatitis A #1	
15 month	Vericella #1	
	Hib (H. influenza type B) #4	
18 month	DTaP (diphtheria, tetanus, and pertussis) #4	
	Hepatitis A #2	
4~6 year	DTaP (diphtheria, tetanus, and pertussis) #5	
	IPV (Polio) #4	
	Vericella #2	
	MMR (Measles, Rubella and Mumps) #2	
9 year-		Human Papillomavirus (HPV)
11-12 year	Tdap (Tetanus, Diphtheria, Pertussis)	
	Meningococcal #1	
16 year	Meningococcal #2	
Every year		Influenza (All people 6 months and older)

※HPV Vaccine: Age 9-14: 2-dose schedule is recommended (0 and 6 month)
Age 15 and older: 3-dose schedule is recommended (0, 1-2 month, 6 month)