



Age	Required	Recommended
Newborn	Hepatitis B #1	Respiratory Syncytial Virus (RSV)*
1 month		
2 month	Hepatitis B #2	Rota#1
	DTaP (diphtheria, tetanus, and pertussis)#1	
	IPV (Polio)#1	
	Hib (H. influenza type B)#1	
	Pneumococcal#1	
4 month	DTaP (diphtheria, tetanus, and pertussis)#2	Rota#2
	IPV (Polio)#2	
	Hib (H. influenza type B)#2	
	Pneumococcal#2	
6 month	Hepatitis B #3	Rota#3
	DTaP (diphtheria, tetanus, and pertussis)#3	
	IPV (Polio)#3	
	Hib (H. influenza type B)#3	
	Pneumococcal#3	
9 month		
12 month	Pneumococcal#4	
	MMR (Measles, Rubella and Mumps)#1	
	Hepatitis A#1	
15 month	Vericella#1	
	Hib (H. influenza type B)#4	
18 month	DTaP (diphtheria, tetanus, and pertussis)#4	
	Hepatitis A#2	
4~6 year	DTaP (diphtheria, tetanus, and pertussis)#5	
	IPV (Polio)#4	
	Vericella#2	
	MMR (Measles, Rubella and Mumps)#2	
9 year-		Human Papillomavirus (HPV)**
11-12 year	Tdap (Tetanus, Diphtheria, Pertussis)	
	Meningococcal #1	
16 year	Meningococcal #2	
Every year		Influenza (All people 6 months and older)

※RSV Vaccine Administration of infant RSV vaccine is recommended during **October through March**.  
An infant RSV antibody is recommended for infants **younger than 8 months of age**

※HPV Vaccine: Age 9-14: 2-dose schedule is recommended (0 and 6 month)  
Age 15 and older: 3-dose schedule is recommended (0, 1-2 month, 6month)